

submitted to *the Astrophysical Journal, Letters*

A NEW ESTIMATION OF MASS ACCUMULATION EFFICIENCY IN HELIUM SHELL FLASHES TOWARD TYPE Ia SUPERNOVA EXPLOSIONS

Mariko Kato

Department of Astronomy, Keio University, Hiyoshi, Kouhoku-ku, Yokohama 223-8521, Japan
e-mail: mariko@educ.cc.keio.ac.jp

and

Izumi Hachisu

Department of Earth Science and Astronomy, College of Arts and Sciences, University of Tokyo,
Komaba, Meguro-ku, Tokyo 153-8902, Japan
e-mail: hachisu@chianti.c.u-tokyo.ac.jp

ABSTRACT

We have calculated the mass accumulation efficiency during helium shell flashes to examine whether or not a carbon-oxygen white dwarf (C+O WD) grows up to the Chandrasekhar mass limit to ignite a Type Ia supernova (SN Ia) explosion. It has been frequently argued that luminous super-soft X-ray sources (SSSs) and symbiotic stars are progenitors of SNe Ia. In such systems, a C+O WD accretes hydrogen-rich matter from a companion and burns hydrogen steadily on its surface. The WD develops a helium layer underneath the hydrogen-rich envelope and undergoes periodic helium shell flashes. Using OPAL opacity, we have reanalyzed a full cycle of helium shell flashes on a $1.3M_{\odot}$ C+O WD and confirmed that the helium envelope of the WD expands to blow a strong wind. A part of the accumulated matter is lost by the wind.

The mass accumulation efficiency in helium shell flashes is estimated as $\eta_{\text{He}} = -0.175 (\log \dot{M} + 5.35)^2 + 1.05$, for $-7.3 < \log \dot{M} < -5.9$, and $\eta_{\text{He}} = 1$ for $-5.9 \leq \log \dot{M} \lesssim -5$, where the mass accretion rate $\dot{M}$ is in units of $M_{\odot} \text{ yr}^{-1}$. In relatively high mass accretion rates as expected in recent SN Ia progenitor models, the mass accumulation efficiency is large enough for C+O WDs to grow to the Chandrasekhar mass, i.e., $\eta_{\text{He}} = 0.9$ for $\log \dot{M} = -6.3$, and $\eta_{\text{He}} = 0.57$ for $\log \dot{M} = -7.0$. The wind velocity ($\sim 1000 \text{ km s}^{-1}$) is much faster than the orbital velocity of the binary ($\lesssim 300 \text{ km s}^{-1}$) and therefore, the wind cannot be accelerated further by the companion's motion. We suggest observational counterparts of helium shell flashes in relation to long term variations in super-soft X-ray fluxes of SSSs and symbiotic stars.

Subject headings: binaries: close — novae, cataclysmic variables — supernova: general
— stars: mass-loss — white dwarfs

1. INTRODUCTION

Type Ia supernovae (SNe Ia) are widely believed to be a thermonuclear explosion event of a white dwarf (WD), although the immediate progenitors have not been identified yet. Among many evolutionary paths toward SNe Ia proposed so far, arguments are now focused on the Chandrasekhar (Ch) mass model vs. the sub-Chandrasekhar (sub-Ch) mass model, and the single degenerate (SD) model vs. the double degenerate (DD) model (e.g., Branch et al. 1995 for a review of SN Ia progenitors). Of these models, the SD & Ch model is the most promising one both from the observational and theoretical points of view (e.g., Nomoto et al. 1994 for a review; Kobayashi et al. 1998).

One example of the SD & Ch models is luminous super-soft X-ray sources (SSSs) in which the WD accretes hydrogen-rich material at relatively high rates ($\dot{M} \gtrsim 1 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$) (van den Heuvel et al. 1992). The WD accretes almost pure helium ($Y = 0.98$) as a results of stable hydrogen shell-burning and experience helium shell flashes. In a strong flash, the envelope expands greatly (e.g., Iben 1982) and may suffer wind mass-loss or a Roche lobe overflow (e.g., Kato, et al. 1989). If a large part of the helium layer will be lost by the wind, the WD hardly grows in mass to the Chandrasekhar mass (Cassisi et al 1998). Thus, it is essentially important to estimate the efficiency of the net mass accumulation onto the C+O core. Kato, Saio, & Hachisu (1989, hereafter KSH89) has first estimated the mass accumulation efficiency in helium shell flashes. They used the old opacity, however, and it should be recalculated with the new opacity.

A new evolutionary path of the SD & Ch scenario has been proposed by Hachisu, Kato, & Nomoto (1996; hereafter HKN96), in which a WD accretes hydrogen-rich matter from a lobe-filling red-giant (WD+RG system). In this model, optically thick winds blow from WDs, which stabilize the mass transfer even if the donor has a deep convective envelope. The WD steadily accretes hydrogen-rich matter from the companion. Its mass eventually reaches the Chandrasekhar mass limit and the WD explodes as an SN Ia. Hachisu, Kato, & Nomoto (1999a, hereafter HKN99) show this WD+RG system is one of the main channels to SNe Ia. Li & van den Heuvel (1997) extended HKN96's model to another type of binary systems consisting of a mass-accreting WD and a lobe-filling, more massive, somewhat evolved main-sequence star (WD+MS system). Hachisu & Kato (1999) further proposed an evolutionary path to binaries consisting of a WD and a somewhat evolved MS star characterized by helium-rich accretion as observed in U Sco. This is another main channel to SNe Ia as shown by Hachisu, Kato, Nomoto, & Umeda (1999b, hereafter HKNU99).

All of these models will experience helium shell flashes followed by optically thick winds during its way to an SN Ia. If most of the helium matter is blown off in the wind, the WD

cannot grow to the Chandrasekhar mass ($M_{\text{Ch}} = 1.38M_{\odot}$). Therefore, once again it is essentially important to evaluate the efficiency of mass accumulation during the wind phase of helium shell flashes in order to reach the definite conclusion of SN Ia progenitors.

Cassisi, Iben & Tornambé (1998, hereafter CIT98) criticized HKN96’s model and concluded that the white dwarf mass hardly increases to M_{Ch} because of heavy mass loss owing to a Roche lobe overflow during helium shell flashes. However, once the optically thick wind occurs during helium shell flashes, the wind velocity is as fast as $\sim 1000 \text{ km s}^{-1}$, much faster than the orbital velocity ($\lesssim 300 \text{ km s}^{-1}$) so that the companion motion hardly affects the mass ejection. The wind matter quickly goes away from the system without any interaction to the orbital motion. Therefore their argument has no physical meaning when a strong wind occurs.

In this *Letter*, we have followed a full cycle of helium shell flashes including the mass loss owing to optically thick winds in order to obtain the mass accumulation efficiency of accreting WDs. We have used the same techniques as in KSH89 but adopted OPAL opacity that has a strong peak around $\log T \text{ (K)} \sim 5.2$, which drives a strong wind as shown by Kato & Hachisu (1994). In §2, we describe our methods, assumptions, and main results of helium shell flashes on a $1.3M_{\odot}$ WD. Discussions follow in §3.

2. MASS LOSS DURING HELIUM SHELL FLASHES

Kato, Saio, & Hachisu (1989, KSH89) adopted two different approaches to follow a full cycle of helium shell flashes: time-dependent hydrostatic calculation and mimicking time-dependent behavior by a sequence of steady-state wind solutions. They combined these two methods, because the time-dependent Henyey code failed when the envelope expands greatly while the steady-state approach works well. We have used the same basic approaches as in KSH89 to follow a full cycle of helium shell flashes on a $1.3M_{\odot}$ WD. The main difference is in the wind solutions, which we replaced by the new solutions recalculated with the updated OPAL opacity (Iglesias & Rogers 1996).

The ignition models are unchanged because OPAL opacity gives essentially the same values as the old opacity does at higher temperatures of $\log T \text{ (K)} > 6$. We have assumed the same white dwarf radius of $1.3M_{\odot}$, i.e., $\log R \text{ (cm)} = 8.513$, and the envelope chemical composition of $Y = 0.48$, $C + O = 0.5$ and $Z = 0.02$ for helium, carbon and oxygen, and heavy elements (including C+O in solar ratio) by weight, respectively. Such a small helium content is expected as follows: in the early stage of a shell flash, the convection spreads over the entire envelope, then descends and disappears before the photosphere reaches the maximum expansion. The ashes of helium burning, i.e., carbon and oxygen, is mixed into the entire envelope to reduce the helium content by about a half on average from the initial value of $Y = 0.98$. Thus, we have assumed the uniform composition with $Y = 0.48$ in the quasi-evolutionary sequence. The convective energy transport is included in both the static and the wind solutions by the mixing length theory with

the mixing-length parameter of 1.5. The convection is, however, inefficient in the wind solutions, because a wind is too fast (supersonic) compared with the convective motion (subsonic). (See Kato & Hachisu 1994 for details.)

Figure 1 depicts evolutionary tracks of one cycle of helium shell flashes. After the onset of a shell flash, the helium layer expands to reach the maximum expansion of the photosphere, the radius of which depends on the ignition mass ΔM_{ig} of the helium layer. We call this *rising phase*. In the rising phase, the optically thick wind occurs when the photospheric temperature decreases to $\log T_{\text{ph}} \sim 5.45$. After the maximum expansion, the photospheric radius decreases with the envelope mass and the star moves blueward in the HR diagram. After the wind stops, the envelope mass further decreases owing to nuclear burning and the star moves downward to come back to the original position. We call this *decay phase*.

For the rising phase, evolutionary tracks are approximated by a sequence of static solutions with a constant envelope mass until the optically thick wind occurs (see KSH89 for details). Three tracks have been calculated with (a) $\Delta M_{\text{ig}} = 4.8 \times 10^{-5} M_{\odot}$, (b) $\Delta M_{\text{ig}} = 8.7 \times 10^{-5} M_{\odot}$, and (c) $\Delta M_{\text{ig}} = 3.5 \times 10^{-4} M_{\odot}$. These ignition masses are realized for the helium accretion rates of (a) $\dot{M}_{\text{acc}} = 1.7 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$, (b) $\dot{M}_{\text{acc}} = 6.7 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$, and (c) $\dot{M}_{\text{acc}} = 1 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$, respectively. The helium accretion rates of (a) and (b) correspond to the largest rates of hydrogen steady shell-burning, $\dot{M}_{\text{acc}} \sim L_{\text{Edd}}/X\varepsilon_H$, in the envelope with $X = 0.35$ and $X = 0.7$, respectively, where $L_{\text{Edd}} \propto 1/(1 + X)$ is the Eddington luminosity for electron scattering and ε_H is the nuclear energy release of hydrogen-burning per unit mass. Thus $\dot{M}_{\text{acc}}$ is proportional to $1/X(1 + X)$ for the hydrogen content X . Such a low hydrogen content of $X = 0.35$ (or $Y = 0.63$) is expected in the helium-rich matter accretion in recurrent nova U Sco (Hachisu & Kato 1999).

The evolutionary track for (a) $\Delta M_{\text{ig}} = 4.8 \times 10^{-5} M_{\odot}$ is very close to the curve for the decay phase (thick curve) and omitted in Figure 1. No optically thick winds occur in this case because the photospheric temperature does not decrease to $\log T_{\text{ph}} \sim 5.45$. At the maximum expansion, the photospheric temperature reaches $\log T_{\text{ph}} = 5.83$ (indicated by arrow a), and the radius $R_{\text{ph}} = 0.022 R_{\odot}$.

In the case of (b) $\Delta M_{\text{ig}} = 8.7 \times 10^{-5} M_{\odot}$, the wind mass-loss begins at $\log T_{\text{ph}} = 5.45$. The maximum expansion is reached at $R_{\text{ph}} \leq 0.70 R_{\odot}$ and $\log T_{\text{ph}} \geq 5.05$ (indicated by arrow b), where the equality stands when we neglect the wind mass loss in the rising phase. The star moves blueward along the thick curve in the decay phase. The envelope mass is decreasing owing to wind and nuclear burning. The optically thick wind ceases at $\log T_{\text{ph}} = 5.45$. In the third case of (c) $\Delta M_{\text{ig}} = 3.5 \times 10^{-4} M_{\odot}$, the wind mass loss begins at $\log T_{\text{ph}} = 5.44$, and the star reaches the maximum expansion at $R_{\text{ph}} \leq 9.1 R_{\odot}$ and $\log T_{\text{ph}} \geq 4.45$ (indicated by arrow c).

These tracks for three different envelope masses merge into one track after the maximum expansion, because the envelope reaches a thermal equilibrium in the decay phase. The more massive the ignition mass is, the more redward the star moves in the HR diagram, that is, the star is traveling a longer track to a lower temperature. If the ignition mass is as large as

$\Delta M_{\text{ig}} \sim 7.7 \times 10^{-4} M_{\odot}$, the star reaches $\log T_{\text{ph}} \geq 4.07$. This envelope mass, $\Delta M_{\text{ig}} = 7.7 \times 10^{-4} M_{\odot}$ corresponds to the ignition mass at $\dot{M}_{\text{acc}} = 4.5 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$.

Figure 2 shows the photospheric temperature T_{ph} , the photospheric radius R_{ph} , and the photospheric wind velocity V_{ph} in the decay phase of helium shell flashes. The arrows indicate the same three ignition masses as in Figure 1. At the maximum expansion, the star reaches these position and then moves blueward as the envelope mass decreases owing to wind mass-loss and nuclear burning. The wind mass-loss stops at $\Delta M = 5.0 \times 10^{-5} M_{\odot}$, marked by a filled circle. The photospheric velocity is as large as $\sim 1000 \text{ km s}^{-1}$, which is much faster than the previous results ($\sim 300 \text{ km s}^{-1}$ in KSH89), because of a strong acceleration driven by the large peak of OPAL opacity.

The newly obtained wind velocities are much faster than the orbital velocities of SN Ia progenitors of the WD+MS or WD+RG systems ($\lesssim 300 \text{ km s}^{-1}$). Therefore, we can neglect effects of the binary motion on the mass ejection during the common envelope phase, because the envelope matter quickly goes away with little interaction with the orbital motion. This implies that neither a Roche lobe overflow nor a common envelope action plays an important role on mass ejection.

Figure 3 shows the total mass decreasing rate of the envelope $\log |d(\Delta M)/dt|$, which is the sum of the wind mass loss rate $\dot{M}_{\text{wind}}$ and the mass decreasing rate owing to nuclear burning $\dot{M}_{\text{nuc}}$ against the envelope mass $\log \Delta M$ for the decay phase of helium shell flashes. The processed matter accumulates at a rate higher than the mass ejection rate for the region of $\log \Delta M < -3.74$, and then a large part of the envelope matter accumulates on the white dwarf. These results are very different from the case of nova outbursts, in which most envelope mass is blown off. It is because the difference is in the nuclear energy release per unit mass between hydrogen-burning and helium-burning: hydrogen burning produces energy ten times larger than that of helium burning, i.e., only a tenth of the hydrogen-rich envelope mass is enough to blow the entire envelope off.

Figure 4 shows the accumulation efficiency defined by the ratio of the processed matter remaining after one cycle of helium shell flash to the ignition mass. The amount of matter lost by the wind in the decay phase is calculated from the rate of wind mass-loss and nuclear burning in Figure 3. This accumulation ratio is well fitted by the solid curve expressed by

$$\eta_{\text{He}} = \begin{cases} 1, & \text{for } -5.9 \leq \log \dot{M} \lesssim -5 \\ -0.175 \left(\log \dot{M} + 5.35 \right)^2 + 1.05, & \text{for } -7.3 < \log \dot{M} < -5.9 \end{cases} \quad (1)$$

where $\dot{M}$ is in units of $M_{\odot} \text{ yr}^{-1}$.

3. DISCUSSIONS

Cassisi et al (1998, CIT98) have followed the evolution of hydrogen-accreting white dwarfs. These WDs are undergoing many cycles of hydrogen shell flashes until a strong helium shell flash

develops to expand the envelope to a red-giant dimension. Based on these calculations with the old opacity, they concluded that a hydrogen-accreting white dwarf will not become an SN Ia, because almost all of the helium envelope is lost from the system by a Roche lobe overflow or a common envelope ejection. CIT98 further denied the possibility of HKN96’s scenario.

In what follows, we criticize CIT98’s argument and support the growth of C+O WDs as expected in HKN96’s and Li & van den Heuvel’s (1997) models. CIT98 calculated only much less massive white dwarfs of $0.516M_{\odot}$ and $0.8M_{\odot}$ to criticize $\sim 1.3M_{\odot}$ model of HKN96. The trends of the envelope expansion and the mass accumulation depend on the WD mass because of the large difference in the surface gravity; nuclear energy release in helium burning is comparable to the gravitational potential energy for massive WDs, i.e., the ratio of nuclear/gravity is ~ 1.0 for a $1.3M_{\odot}$ WD, whereas it is as large as 3.9 for a $0.8M_{\odot}$ WD, and 8.9 for a $0.5M_{\odot}$ WD. For massive WDs, therefore, a large part of helium is consumed to produce nuclear energy before the envelope expands against the gravity. It reduces the helium content of the envelope. This low helium content in the envelope increases the burning speed in mass coordinates as understood from $\dot{M}_{\text{nuc}} \approx L_{\text{Edd}}/Y\varepsilon_{\text{He}}$, where ε_{He} is the nuclear energy release per unit mass for helium burning. Thus, CIT98’s criticism is inappropriate in the sense that their results for less massive WDs cannot be directly applied to very massive WDs.

Once the optically thick wind occurs, a Roche lobe overflow or a common envelope ejection discussed in CIT98 becomes physically meaningless, because the wind velocity $\sim 1000 \text{ km s}^{-1}$ is much faster than the orbital velocity of the companion ($\lesssim 300 \text{ km s}^{-1}$) of typical SN Ia progenitors (WD+MS or WD+RG model). The wind goes away quickly without any effective acceleration by the companion. After the wind stops, the photospheric radius ($\lesssim 0.1R_{\odot}$) is much smaller than the Roche lobe ($\gtrsim 1R_{\odot}$). Therefore, neither a Roche lobe overflow nor a common envelope ejection works in our binary system.

We have ignored hydrogen-rich envelope above the helium layer. An accreting $1.3 M_{\odot}$ WD has a hydrogen-rich envelope of $2 \times 10^{-7} - 3 \times 10^{-6}M_{\odot}$ above the helium layer, depending on the mass transfer rate, chemical composition, and the other details of binary models. This is much smaller ($\lesssim 0.01$) than the mass of the helium layer at ignition and we have neglected its contribution to the mass accumulation efficiency. The hydrogen-rich envelope will be quickly blown off at a very early stage of helium shell flashes.

Helium shell flashes may be observed as transient super-soft X-ray sources. For high mass accretion rates as expected in SN Ia progenitors, shell flashes are so weak that the surface temperature never decrease to less than $\sim 1 \times 10^5 \text{ K}$. Therefore, it will be a bright super-soft X-ray phenomenon with a faint optical counterpart. For a $1.3M_{\odot}$ WD, the bright SSS stage lasts $\sim 1 - 3 \text{ yrs}$ with recurrence period of a few tens to several tens of years. For more massive WDs just before explosions, it lasts several months with the recurrence period of several years.

This research has been supported in part by the Grant-in-Aid for Scientific Research

(08640321, 09640325) of the Japanese Ministry of Education, Science, Culture, and Sports.

REFERENCES

- Branch, D., Livio, M., Yungelson, L.R., Boffi, F.R., Baron, E. 1995, *PASP*, 107, 717
- Cassisi, S., Iben, I. Jr. & Tornambé, A., 1998, *ApJ*, 496, 376 (CIT98)
- Hachisu, I. & Kato, M. 1999, *ApJ*, submitted
- Hachisu, I., Kato, M., Nomoto, K. 1996, *ApJ*, 470, L97 (HKN96)
- Hachisu, I., Kato, M., Nomoto, K. 1999a, *ApJ*, submitted (HKN99)
- Hachisu, I., Kato, M., Nomoto, K. & Umeda, H. 1999b, *ApJ*, submitted (HKNU99)
- Iben, I. Jr. 1982, *ApJ*, 259, 244
- Iglesias, C. A., Rogers, F. 1996, *ApJ*, 464, 943
- Kato, M., Hachisu, I., 1994, *ApJ*, 437, 802
- Kato, M., Saio, H., & Hachisu, I., 1989, *ApJ*, 340, 509 (KSH89)
- Kobayashi, C., Tsujimoto, T., Nomoto, K., Hachisu, I., & Kato, M. 1998, *ApJ*, 503, L155
- Li, X.-D., van den Heuvel, E. P. J. 1997, *A&A*, 322, L9
- Nomoto, K., Yamaoka, H., Shigeyama, T., Kumagai, S., Tsujimoto, T. 1994, in *Supernovae (Les Houches, Session LIV)*, ed. S. Bludman et al. (Elsevier Sci. Publ.), p. 199
- van den Heuvel, E. P. J., Bhattacharya, D., Nomoto, K., & Rappaport, S. 1992, *A&A*, 262, 97




Fig. 1.— Evolutionary tracks of one cycle of helium shell flash on a $1.3 M_{\odot}$ white dwarf. Two tracks of the rising phase are shown by thin solid curves: the ignition mass is (b) $\Delta M_{\text{ig}} = 8.7 \times 10^{-5} M_{\odot}$ (upper), and (c) $3.5 \times 10^{-4} M_{\odot}$ (lower). The maximum expansion of the photosphere is indicated by an arrow for three ignition masses of (a) $\Delta M_{\text{ig}} = 4.8 \times 10^{-5} M_{\odot}$, (b) $\Delta M_{\text{ig}} = 8.7 \times 10^{-5} M_{\odot}$, and (c) $\Delta M_{\text{ig}} = 3.5 \times 10^{-4} M_{\odot}$. The decay phases for these three ignition masses are merged into one line (thick solid). On the decay phase, the optically thick wind ceases at the point marked by the cross. The decay phase is extended to $\log T = 4.07$ (short-dashed) for a strong helium shell flash with a large initial envelope mass $\Delta M_{\text{ig}} = 7.7 \times 10^{-4} M_{\odot}$.




Fig. 2.— Photospheric radius R_{ph} (cm), temperature T_{ph} (K), and wind velocity V_{ph} (cm s^{-1}), are plotted against the envelope mass ΔM for the decay phase of helium shell flashes. Time runs from right to left. The wind mass-loss stops at $\Delta M = 5.0 \times 10^{-5} M_{\odot}$ denoted by the dots. The three ignition masses shown in Fig. 1 are indicated by the arrows.




Fig. 3.— Wind mass-loss rate $\dot{M}_{\text{wind}}$ (dashed), the mass decreasing rate owing to helium nuclear burning $\dot{M}_{\text{nuc}}$ (dot-dashed), and the total envelope mass decreasing rate $\dot{M}_{\text{wind}} + \dot{M}_{\text{nuc}}$ (solid) are plotted against the envelope mass ΔM . The ignition mass of three cases are shown by the arrows. Time runs from right to left.




Fig. 4.— Mass accumulation efficiency η_{He} , the ratio of the mass accreted after one cycle of helium shell flash to the ignition mass, is plotted against the helium accretion rate (filled circles). The solid curve is an empirical formula given by equation (1). The three mass accretion rates are indicated by the arrows, which are corresponding to the three ignition masses shown in Fig. 1.